

Stability of Spreading Processes in Networks with Non-Markovian Transmission and Recovery

Masaki Ogura, *Member, IEEE*, Victor M. Preciado, *Member, IEEE*

Abstract—Although viral spreading processes taking place in networks are commonly analyzed using Markovian models in which both the *transmission* and the *recovery* times follow exponential distributions, empirical studies show that, in most real scenarios, the distribution of these times are far from exponential. To overcome this limitation, we first introduce a generalized spreading model that allows transmission and recovery times to follow arbitrary distributions within a given accuracy. In this context, we derive conditions for the generalized spreading model to converge exponentially fast towards the infection-free equilibrium with a given rate (in other words, to eradicate the viral spread) without relying on mean-field approximations. Based on our results, we illustrate how the particular shape of the transmission/recovery distribution heavily influences the convergence rate. Therefore, modeling non-exponential transmission/recovery times observed in realistic spreading processes via exponential distributions can induce significant errors in the stability analysis of the spreading dynamics.

I. INTRODUCTION

Understanding the dynamics of spreading processes in complex networks is a challenging problem with a wide range of practical applications in epidemiology and public health [2], or information propagation in social networks [10]. During the last decade, significant progress has been made towards understanding the relationship between the topology of a network and the dynamics of spreading processes taking place over the network (see [11] for a recent survey). A common approach to investigate this relationship is by modeling spreading processes using networked Markov processes, such as the networked susceptible-infected-susceptible (SIS) model [17]. Based on these Markovian models, it is then possible to find an explicit relationship between epidemic thresholds and network eigenvalues in static topologies [7], [17], as well as in temporal [12], and adaptive [13] networks.

A consequence of using Markovian models in the analysis of spreading processes is that both *transmission times* (i.e., the time it takes for an infection to be transmitted from an infected node to one of its neighbors) and *recovery times* (i.e., the time it takes for an infected node to recover) follow exponential distributions. However, empirical studies show that, in most real networks, the distribution of transmission and recovery times are far from exponential. For example, the transmission time of messages in Twitter, or news in other social media outlets, follows (approximately) a log-normal distribution [10].

In the context of human contact networks, the transmission of immunodeficiency viruses [3], or the time it takes to recover from the influenza virus [16] are clearly non-exponential.

Since realistic transmission and recovery times usually follow non-exponential distributions, it is of practical importance to understand the role of these distributions on the dynamics of the spread. In this direction, the authors in [18] illustrated via numerical simulations that non-exponential transmission times can have a substantial effect on the speed of spreading. An approximate analysis of spreading processes with general transmission and recovery times using mean-field approximations was proposed in [4]. In [9], an analytically solvable (although rather simplistic) spreading model with non-exponential transmission times was proposed. Using moment-closure techniques, the authors in [15] analyzed spreading processes with non-exponential transmission and recovery times in special network topologies (in particular, triangles and chains of length 2).

In this paper, we propose a tractable approach to analyze spreading processes over arbitrary networks with general transmission and recovery times. In this direction, we first introduce the *generalized networked SIS* (GeNeSIS) model, which allows for transmission and recovery times following arbitrary phase-type distributions (see, e.g., [1]). Since phase-type distributions form a dense family in the space of positive-valued distributions [6], the GeNeSIS model allows to theoretically analyze arbitrary transmission and recovery times within an arbitrary accuracy [1]. We are particularly interested in finding conditions for the dynamics of the spread to converge exponentially fast towards the infection-free equilibrium with a given rate; in other words, to eradicate the viral spreading process. In contrast with existing results [4], [15], we provide an analysis of the dynamics over *arbitrary* networks with general infection and recovery distributions (without relying on mean-field approximations). The key tool used in our derivations is a vectorial representation of phase-type distributions, which we use to derive conditions for exponential stability of the infection-free equilibrium in the stochastic dynamics of the GeNeSIS model. Based on our theoretical results, we show that modeling non-exponential transmission/recovery times using exponential distributions induces significant errors in the dynamics of spreading processes, as previously observed via numerical simulations.

This paper is organized as follows. In Section II, we introduce elements of graph theory and stochastic differential equations with Poisson jumps. In Section III, we describe a generalized model of spreading processes over networks with arbitrary transmission and recovery times. In Section IV,

The authors are with the Department of Electrical and Systems Engineering, University of Pennsylvania, Philadelphia, PA 19014, USA. Email: {ogura,preciado}@seas.upenn.edu

This work was supported in part by the NSF under grants CNS-1302222 and IIS-1447470.

we provide a vectorial representation of the GeNeSIS model, which we use in Section V to derive stability conditions of its stochastic dynamics. We validate the effectiveness of our results via numerical simulations in Section VI, where we also illustrate the significant effect of non-exponential transmission/recovery times in the dynamics of the spread.

II. MATHEMATICAL PRELIMINARIES

Let $\mathbb{R}$ and $\mathbb{N}$ denote the set of real numbers and positive integers, respectively. For a positive integer n , define $[n] = \{1, \dots, n\}$. For a real function f , let $f(t^-)$ denote the limit of f from the left at time t . We let I_n and O_n denote the $n \times n$ identity and zero matrices. By $\mathbb{1}_p$ and $\mathbb{0}_p$, we denote the p -dimensional vectors whose entries are all ones and zeros, respectively. A real matrix A (or a vector as its special case) is said to be nonnegative, denoted by $A \geq 0$, if A is nonnegative entry-wise. The notation $A \leq 0$ is understood in the obvious manner. For a square matrix A , the maximum real part of its eigenvalues, called the *spectral abscissa* of A , is denoted by $\eta(A)$. We say that A is Metzler if the off-diagonal entries of A are all non-negative. It is easy to see that, if A is Metzler, then $e^{At} \geq 0$ for every $t \geq 0$. The Kronecker product [8] of two matrices A and B is denoted by $A \otimes B$, and the Kronecker sum of two square matrices $A \in \mathbb{R}^{p \times p}$ and $B \in \mathbb{R}^{q \times q}$ is defined by $A \oplus B = A \otimes I_q + I_p \otimes B$. Given a collection of n matrices $A_1, \dots, A_n$ having the same number of columns, the matrix obtained by stacking the matrices in vertical (A_1 on top) is denoted by $\text{col}(A_1, \dots, A_n)$.

An undirected graph is a pair $\mathcal{G} = (\mathcal{V}, \mathcal{E})$, where $\mathcal{V} = \{1, \dots, n\}$ is the set of nodes, and $\mathcal{E} \subset \mathcal{V} \times \mathcal{V}$ is the set of edges, consisting of distinct and unordered pairs $\{i, j\}$ for $i, j \in \mathcal{V}$. We say that a node i is a neighbor of j (or that i and j are adjacent) if $\{i, j\} \in \mathcal{E}$. The set of neighbors of node i is denoted by $\mathcal{N}_i$. The adjacency matrix of $\mathcal{G}$ is defined as the $n \times n$ matrix whose (i, j) -th entry is 1 if and only if nodes i and j are adjacent, 0 otherwise.

Let $(\Omega, \mathcal{M}, P)$ be a probability space. The expectation of a random variable is denoted by $E[\cdot]$. A Poisson counter [5, Chapter 4] of rate $\lambda > 0$ is denoted by N_λ . In this paper, we extensively use a specific class of stochastic differential equations with Poisson jumps, described below. For each $i \in [m]$, let $f_i: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function, N_{λ_i} be a Poisson counter, and κ_i be a continuous-time real stochastic process defined over the probability space Ω such that $\{\kappa_i(t)\}_{t \geq 0}$ are independent and identically distributed. All the above stochastic processes are assumed to be independent of each other. Then, we say that a real and right-continuous function x is a solution of the stochastic differential equation

$$dx = \sum_{i=1}^m f_i(x(t), \kappa_i(t)) dN_{\lambda_i}, \quad (1)$$

if x is constant on any interval where none of the counters $N_{\lambda_1}, \dots, N_{\lambda_m}$ jumps, and $x(t) = x(t^-) + f_i(x(t^-), \kappa_i(t))$ when N_{λ_i} jumps at time t . This definition can be naturally extended to the vector case. Below, we present two lemmas for this class of stochastic differential equations. The first lemma states a version of Itô's formula:

Lemma II.1. Assume that x is a solution of (1). Let g be a real continuous function. Then, $y(t) = g(x(t))$ is a solution of the stochastic differential equation $dy = \sum_{i=1}^m [g(x(t) + f_i(x(t), \kappa_i(t))) - g(x(t))] dN_{\lambda_i}$, i.e., y satisfies $y(t) = y(t^-) + g(x(t^-) + f_i(x(t^-), \kappa_i(t))) - g(x(t^-))$, if the Poisson counter N_{λ_i} jumps at time t and is constant over any interval in which none of the counters $N_{\lambda_1}, \dots, N_{\lambda_m}$ jumps.

Proof. Assume that the counter N_{λ_i} jumps at time t . Since x is the solution of the stochastic differential equation (1), it follows that $y(t) - y(t^-) = g(x(t^-) + f_i(x(t^-), \kappa_i(t))) - g(x(t^-))$, as desired. $\square$

We also state the following lemma concerning the expectation of the solution to the stochastic differential equation (1). The proof is omitted because it follows from an easy application of Dynkin's formula.

Lemma II.2. Assume that x is a solution of (1). If the functions $f_1, \dots, f_m$ are affine with respect to the second variable, then $dE[x(t)]/dt = \sum_{i=1}^m E[f_i(x(t), E[\kappa_i(t)])] \lambda_i$.

III. SIS MODEL WITH GENERAL TRANSMISSION AND RECOVERY TIMES

The aim of this section is to introduce the generalized networked susceptible-infected-susceptible (GeNeSIS) model, which will allow us to analyze the effect of arbitrary transmission and recovery times in the spreading dynamics.

A. Generalized Networked SIS Model

Consider a network of n nodes¹ connected via an undirected graph $\mathcal{G} = (\mathcal{V}, \mathcal{E})$. In this model, the state of a node $i \in \mathcal{V}$ is described by a $\{0, 1\}$ -valued continuous-time stochastic process, denoted by $z_i = \{z_i(t)\}_{t \in \mathbb{R}}$. We say that node i is *susceptible* (respectively, *infected*) at time t if $z_i(t) = 0$ (respectively, $z_i(t) = 1$). We assume that the function z_i is continuous from the right for all $i \in [n]$. Under this assumption, we say that node i *becomes infected* (respectively, *becomes susceptible*) at time t if $z_i(t^-) = 0$ and $z_i(t) = 1$ (respectively, $z_i(t^-) = 1$ and $z_i(t) = 0$). It is assumed that all nodes are susceptible before time $t = 0$, i.e., $z_i(t) = 0$ for $t < 0$.

Using this notation, we introduce the GeNeSIS model as follows.

Definition III.1. We say that the family $z = \{z_i\}_{i \in [n]}$ of stochastic processes is a *generalized networked susceptible-infected-susceptible model* (GeNeSIS model, for short) if there exist a subset $\Lambda \subset [n]$, as well as random variables $0 = \tau_0^{ji}(t) < \tau_1^{ji}(t) < \dots$, and $\rho^i(t) > 0$ satisfying the following conditions for all $i \in [n]$, $j \in \mathcal{N}_i$, and $t \geq 0$:

- Node i becomes infected at time $t = 0$ if and only if $i \in \Lambda$, i.e., Λ is the initially infected subset.
- Assume that node i becomes infected at time t . Then, node i remains infected during the time interval $[t, t + \rho^i(t))$ and becomes susceptible at time $t + \rho^i(t)$, i.e., the random variable $\rho^i(t)$ is the *recovery time* of node i ;

¹Without loss of generality, we assume that each node has at least one neighbor.

- c) During the interval $[t, t + \rho^i(t))$, the infected node i attempts to infect node $j \in \mathcal{N}_i$ at times $\{t + \tau_k^{ji}(t)\}_{k \in \mathbb{N}}$. If node j is susceptible at time $t + \tau_k^{ji}(t)$ for any $k \in \mathbb{N}$, then it becomes infected.

Remark III.2. We call the random increments $\{\tau_k^{ji}(t) - \tau_{k-1}^{ji}(t)\}_{j \in \mathcal{N}_i, t \geq 0, k \in \mathbb{N}}$ the *transmission times* of node i , since the difference $\tau_k^{ji}(t) - \tau_{k-1}^{ji}(t)$ represents the time between infection attempts from an infected node i towards a neighboring node j . Note that, when all the recovery and transmission times follow exponential distributions, the GeNeSIS model recovers the standard networked SIS model studied in the literature [11], [17].

Notice that the origin (i.e., $z_i = 0$ for all $i \in \mathcal{V}$) is an absorbing state of the GeNeSIS dynamics. In what follows, we will refer to the origin as the *infection-free* equilibrium. The aim of this paper is to analyze the dynamics of the generalized SIS model according to the following stability criterion:

Definition III.3. For a given $\lambda > 0$, we say that the infection-free equilibrium of the GeNeSIS model is *exponentially mean stable with decay rate λ* if, for every $\varepsilon > 0$, there exists a $C > 0$ such that $E[z_i(t)] \leq Ce^{-(\lambda - \varepsilon)t}$ for all $t \geq 0$, $i \in [n]$, and $\Lambda \subset [n]$.

B. Phase-type Transmission and Recovery Times

In this paper, we consider the GeNeSIS model with transmission and recovery times following phase-type distributions [1]. In what follows, we briefly describe this wide class of probability distributions. Consider a time-homogeneous Markov process x in continuous-time with $p + 1$ ($p \in \mathbb{N}$) states (also called *phases*) such that the states $1, \dots, p$ are transient and the state $p + 1$ is absorbing. The infinitesimal generator of the process is then necessarily of the form $\begin{bmatrix} T & b \\ 0 & 0 \end{bmatrix}$ and $b = -T\mathbf{1}_p$, where $T \in \mathbb{R}^{p \times p}$ is an invertible Metzler matrix with non-positive row-sums. Let $\text{col}(\phi, 0) \in \mathbb{R}^{p+1}$ ($\phi \in \mathbb{R}^p$) denote the initial distribution of the Markov process x , i.e., $P(x(0) = m) = \phi_m$ for $m \in [p]$ and $P(x(0) = p + 1) = 0$. Then, the time to absorption into the state $p + 1$ is a random variable following a *phase-type distribution*, which we denote by the pair (ϕ, T) . In the rest of the paper, we make the following assumption on the distribution of (random) transmission and recovery times in the GeNeSIS model:

Assumption III.4. Transmission and recovery times of all nodes follow phase-type distributions (ϕ, T) and (ψ, R) , respectively.

Remark III.5. It is known that the set of phase-type distributions is dense in the set of positive-valued distributions [6]. Therefore, it is possible to approximate any arbitrary distribution using a phase-type distribution within any given accuracy. Moreover, there are efficient numerical algorithms to find the parameters of this approximating phase-type distribution [1].

IV. VECTORIAL REPRESENTATIONS

The aim of this section is to introduce a useful vectorial representation of the GeNeSIS model under Assumption III.4.

We start our exposition by providing a vectorial representation of an arbitrary phase-type distribution (Subsection IV-A), and then present a vectorial representation of the GeNeSIS model (Subsection IV-B).

A. Vectorial Representation of Phase-Type Distributions

In what follows, we use the following notation: For $m, m' \in [p]$, let $E_{mm'}$ denote the $p \times p$ matrix whose entries are all zeros, except for its (m, m') -th entry being one. Also, let e_m denote the m -th vector of the canonical basis in $\mathbb{R}^p$ (i.e., all the entries of e_m are zeros, except for the m -th entry being one). Finally, given a probability distribution ϕ on $[p]$, we say that an $\mathbb{R}^p$ -valued random variable x follows the distribution ϕ , denoted by $x \sim \phi$, if $P(x = e_m) = \phi_m$ for every $m \in [p]$.

In the following proposition, we prove that a random variable following a phase-type distribution can be represented using the *exit time* (see [14, p. 117]) of a particular stochastic differential equation:

Proposition IV.1. Let (ϕ, T) be a phase-type distribution. Consider the $\mathbb{R}^p$ -valued stochastic differential equation $dx = \sum_{m, m'=1}^p (E_{m'm} - E_{mm})x dN_{T_{mm'}} - \sum_{m=1}^p E_{mm}x dN_{b_m}$, with random initial condition $x(0) \sim \phi$. Then, the random variable

$$\begin{aligned} \rho &= \min\{t > 0: x(t) = 0\} \\ &= \min\{t > 0: \exists m \in [p] \text{ such that } x(t^-) = e_m, \\ &\quad \text{and } N_{b_m} \text{ jumps at time } t\} \end{aligned} \quad (2)$$

follows (ϕ, T) .

Proof. From the stochastic differential equation in the proposition, we can see that its solution x is a Markov process having the state space $\{e_1, \dots, e_p, 0\} \subset \mathbb{R}^p$, the infinitesimal generator $\begin{bmatrix} T & b \\ 0 & 0 \end{bmatrix}$, and the unique absorbing state 0. Therefore, since $x(0)$ follows the probability distribution ϕ , the random variable ρ follows the phase-type distribution (ϕ, T) . Furthermore, the latter equality in (2) easily follows from the second term in the right-hand side of the differential equation. This completes the proof. $\square$

Based on Proposition IV.1, we now provide a vectorial representation of *renewal sequences* (see, e.g., [5, Chapter 9]) whose inter-renewal times follow a phase-type distribution:

Proposition IV.2. Let (ϕ, T) be a phase-type distribution. Let $e_\phi = \{e_\phi(t)\}_{t \geq 0}$ be a stochastic process such that $e_\phi(t)$ follows the distribution ϕ independently for each $t \geq 0$. Consider the $\mathbb{R}^p$ -valued stochastic differential equation $dx = \sum_{m, m'=1}^p (E_{m'm} - E_{mm})x dN_{T_{mm'}} + \sum_{m=1}^p (e_\phi(t)e_m^\top - E_{mm})x dN_{b_m}$ with a random initial state $x(0)$ following the distribution ϕ . Define $\tau_0 = 0$ and let $0 < \tau_1 < \tau_2 < \dots$ be the (random) times at which $x(t^-) = e_m$ and the counter N_{b_m} jumps for some $m \in [p]$. Then, the increments $\{\tau_k - \tau_{k-1}\}_{k \in \mathbb{N}}$ are independent and identically distributed random variables following (ϕ, T) .

Proof. Since the stochastic differential equation given in the proposition is equivalent to the one in Proposition IV.1 on the time interval $[0, \tau_1)$, the random variable τ_1 has the same probability distribution as the random variable ρ defined in (2) and, therefore, follows the phase-type distribution (ϕ, T) by

Proposition IV.1. Furthermore, by the definition of τ_1 , there exists an $m \in [p]$ such that $x(\tau_1) = (e_\phi(\tau_1)e_m^\top - E_{mm})x(\tau_1^-) + x(\tau_1^-) = e_\phi(\tau_1)$, which follows ϕ . Therefore, by the same argument as above, we see that the random increment $\tau_2 - \tau_1$ also follows (ϕ, T) . An induction completes the proof. $\square$

B. Vectorial Representation of the Generalized SIS Model

In this subsection, we use Propositions IV.1 and IV.2 to provide a vectorial representation of the GeNeSIS model under Assumption III.4. Let $A = [a_{ij}]_{i,j}$ be the adjacency matrix of the graph $\mathcal{G}$. Define the vectors

$$b = -T\mathbb{1}_p \in \mathbb{R}^p, \quad d = -R\mathbb{1}_q \in \mathbb{R}^q. \quad (3)$$

For $\ell, \ell' \in [q]$, let $F_{\ell\ell'}$ denote the $q \times q$ matrix whose entries are all zeros, except for its (ℓ, ℓ') -th entry being one. Also, let f_ℓ denote the ℓ -th vector in the canonical basis of $\mathbb{R}^q$. Finally, for $i, j \in [n]$ and $\gamma > 0$, we let N_{λ}^{ij} and N_{λ}^i denote independent Poisson counters with rate λ . The next theorem is the first main result of this paper:

Theorem IV.3. *For each $i \in [n]$ and $j \in \mathcal{N}_i$, let $e_\phi^{ji} = \{e_\phi^{ji}(t)\}_{t \geq 0}$ and $f_\psi^i = \{f_\psi^i(t)\}_{t \geq 0}$ be the stochastic processes such that $e_\phi^{ji}(t) \sim \phi$ and $f_\psi^i(t) \sim \psi$, independently. Let x^{ji} and y^i be, respectively, the $\mathbb{R}^p$ - and $\mathbb{R}^q$ -valued stochastic processes satisfying the following stochastic differential equations:*

$$dx^{ji} = \sum_{m,m'=1}^p (E_{m'm} - E_{mm})x^{ji} dN_{T_{mm'}}^{ji} + \sum_{m=1}^p (e_\phi^{ji}e_m^\top - E_{mm})x^{ji} dN_{b_m}^{ji} - x^{ji} \sum_{\ell=1}^q y_\ell^i dN_{d_\ell}^i + e_\phi^{ji}(1 - \mathbb{1}_q^\top y^i) \sum_{k=1}^n a_{ik} \sum_{m=1}^p x_m^{ik} dN_{b_m}^{ik}, \quad (4)$$

$$dy^i = \sum_{\ell,\ell'=1}^q (F_{\ell\ell'} - F_{\ell\ell})y^i dN_{R_{\ell\ell'}}^i - y^i \sum_{\ell=1}^q y_\ell^i dN_{d_\ell}^i + f_\psi^i(1 - \mathbb{1}_q^\top y^i) \sum_{k=1}^n a_{ik} \sum_{m=1}^p x_m^{ik} dN_{b_m}^{ik}, \quad (5)$$

where, for an initially infected subset $\Lambda \subset [n]$, the initial conditions satisfy $x^{ji}(0) \sim \phi$ and $y^i(0) \sim \psi$ if $i \in \Lambda$, and $x^{ji}(0) = 0_p$ and $y^i(0) = 0_q$ otherwise. Then, the generalized networked SIS model in Definition III.1 with transmission and recovery times following, respectively, phase-type distributions (ϕ, T) and (ψ, R) can be equivalently described as the family of stochastic processes $z = \{z_i\}_{i=1}^n$, where $z_i(t) = \mathbb{1}_q^\top y_i(t)$ for all $t \geq 0$ and $i \in [n]$.

Before we provide a proof of Theorem IV.3, we start from the following result:

Lemma IV.4. *The following statements are true for all $i \in [n]$, $j \in \mathcal{N}_i$, and $t \geq 0$: 1) $\mathbb{1}_p^\top x^{ji}(t) = \mathbb{1}_q^\top y^i(t)$; 2) $x^{ji}(t) \in \{0_p, e_1, \dots, e_p\}$; and 3) $y^i(t) \in \{0_q, f_1, \dots, f_q\}$.*

Proof. To prove the first statement, fix $i \in [n]$ and $j \in \mathcal{N}_i$, and let $\varepsilon = \mathbb{1}_p^\top x^{ji} - \mathbb{1}_q^\top y^i$. From (4) and (5), we can show that $d\varepsilon = \varepsilon \sum_{\ell=1}^q y_\ell^i dN_{d_\ell}^i$. This equation implies that ε is constant over $[0, \infty)$ because $\varepsilon(0) = 0$. Therefore, we have $\varepsilon(t) = \varepsilon(0) = 0$ for every $t \geq 0$, completing the proof of the first statement.

Let us prove the second statement. Notice that x^{ji} and y^i are piecewise constant since they are the solutions of the stochastic differential equations (4) and (5). Moreover, their values can change only when one of the Poisson counters in the stochastic differential equations jumps. Finally, the initial conditions in Theorem IV.3 shows $x^{ji}(0) \in U = \{0_p, e_1, \dots, e_p\}$. Therefore, it is sufficient to show that the jump of any Poisson counter leaves the set U invariant. Let $t > 0$ be arbitrary and assume $x^{ji}(t^-) \in U$. The stochastic differential equations (4) and (5) have the following five different types of Poisson counters: $N_{T_{mm'}}^{ji}$, $N_{R_{\ell\ell'}}^i$, $N_{b_m}^{ji}$, $N_{b_m}^{ik}$, and $N_{d_\ell}^i$. We now show below that any of these counters leave U invariant. Case 1) When $N_{T_{mm'}}^{ji}$ jumps at time t , we have $x^{ji}(t) = x^{ji}(t^-) + (E_{m'm} - E_{mm})x^{ji}(t^-) = e_{m'}$ if $x^{ji}(t^-) = e_m$, and $x^{ji}(t) = x^{ji}(t^-)$ otherwise, which implies $x^{ji}(t) \in U$. Case 2) When $N_{R_{\ell\ell'}}^i$ jumps at time t , clearly $x^{ji}(t) = x^{ji}(t^-) \in U$. Case 3) When $N_{b_m}^{ji}$ jumps at time t , we can show $x^{ji}(t) \in U$ because $x^{ji}(t) = x^{ji}(t^-) + (e_\phi^{ji}(t)e_m^\top - E_{mm})x^{ji}(t^-)$ if $x_m^{ji}(t^-) = 1$, and $x^{ji}(t) = x^{ji}(t^-)$ otherwise. Case 4) When $N_{b_m}^{ik}$ jumps at time t , we have that $x^{ji}(t) = x^{ji}(t^-) + e_\phi^{ji}(t)(1 - \mathbb{1}_q^\top y^i(t^-))a_{ik}x_m^{ik}(t^-) = x^{ji}(t^-)$ if $\mathbb{1}_q^\top y^i(t^-) = 1$, $a_{ik} = 0$, or $x_m^{ik}(t^-) = 0$, and $x^{ji}(t) = e_\phi^{ji}(t)$ otherwise. Therefore, $x^{ji}(t) \in U$. Case 5) When $N_{d_\ell}^i$ jumps at time t , we have $x^{ji}(t) = x^{ji}(t^-) - x^{ji}(t^-)y_\ell^i(t^-) = 0$ if $y_\ell^i(t^-) = 1$, and $x^{ji}(t) = x^{ji}(t^-)$ otherwise. Therefore, $x^{ji}(t) \in U$. We have thus proved that the set U is left invariant by any of the counters.

The proof of the third statement is similar to the proof of the second statement and, therefore, is omitted. $\square$

From the proof of Lemma IV.4, the following corollary immediately follows.

Corollary IV.5. *The following statements are true for every $i \in [n]$, $j \in \mathcal{N}_i$, and $t \geq 0$:*

- 1) *Node i attempts to infect node j at time t , if and only if, $x_m^{ji}(t^-) = 1$ and $N_{b_m}^{ji}$ jumps at time t for some $m \in [p]$.*
- 2) *Node i becomes susceptible at time t , if and only if, $y_\ell^i(t^-) = 1$ and $N_{d_\ell}^i$ jumps at time t for some $\ell \in [q]$.*

Proof. From the proof of Lemma IV.4, the value of $z^i = \mathbb{1}_p^\top x^{ji} = \mathbb{1}_q^\top y^i$ can change from zero to one only in Case 4), given in the proof. From the equations in Case 4), we see that node i becomes infected at time $t \geq 0$, if and only if, $z_i(t^-) = 0$, node i is adjacent to node k , $x_m^{ik}(t^-) = 1$, and $N_{b_m}^{ik}$ jumps at time t for some $m \in [p]$. Therefore, node k attempts to infect node i at time t if and only if $x_m^{ik}(t^-) = 1$ and $N_{b_m}^{ik}$ jumps at time t for some $m \in [p]$, as desired. Similarly, we see that the value of z^i can change from one to zero only in Case 5). The equations in Case 5) imply that node i becomes susceptible at time t if and only if $y_\ell^i(t^-) = 1$ and the counter $N_{d_\ell}^i$ jumps at time t for some $\ell \in [q]$, showing the second assertion of the corollary. $\square$

Now we are ready to prove our first main result, namely, Theorem IV.3.

Proof of Theorem IV.3. We prove that the family of stochastic processes $z = \{z_i\}_{i=1}^n$, defined as the solution of (4) and (5), satisfies items a)-c) in Definition III.1, as well as Assumption III.4. Item a) is true by the given initial conditions. Let

us now prove item b). Assume that node i becomes infected at time $t \geq 0$; hence, we have that either $t = 0$ or $t > 0$. In this proof, we only consider the case $t = 0$, as the other case can be proved in a similar way. Let ρ be the earliest time at which $y^i(\rho) = 0_q$. Then, on the time interval $[0, \rho)$, the differential equation (5) is equivalent to $dy^i = \sum_{\ell, \ell'=1}^q (F_{\ell'\ell} - F_{\ell\ell'}) y^i dN_{R_{\ell\ell'}}^i - \sum_{\ell=1}^q F_{\ell\ell} y^i dN_{d_\ell}^i$, since $y^i y_\ell^i = F_{\ell\ell} y^i$. Moreover, the vector $y^i(0)$ follows the distribution ψ . Therefore, by Proposition IV.1, we see that ρ follows the phase-type distribution (ψ, R) , proving item b) in Definition III.1 under Assumption III.4 for $t = 0$.

Let us finally prove that the family of stochastic processes $z = \{z_i\}_{i=1}^n$ satisfies item c) in Definition III.1 under Assumption III.4. Assume that node i becomes infected at time t , and, again, consider only the case $t = 0$. Let $\tau_1 < \tau_2 < \dots$ be the times at which node i attempts to infect node j . We need to show that the differences $\tau_1 - 0 = \tau_1, \tau_2 - \tau_1, \dots$ are independent and identically distributed random variables following the phase-type distribution (ϕ, T) . Notice that $x^{ji}(t) \neq 0$ until node i becomes susceptible, which occurs at the earliest time $t > 0$ at which $y_\ell^i(t^-) = 1$ and the counter $N_{d_\ell}^i$ jumps for some $\ell \in [q]$ (by Corollary IV.5). Therefore, until the recovery of node i , the stochastic differential equation (4) is equivalent to $dx^{ji} = \sum_{m, m'=1}^p (E_{m'm} - E_{mm}) x^{ji} dN_{T_{mm'}}^{ji} + \sum_{m=1}^p (e_\phi^j e_m^\top - E_{mm}) x^{ji} dN_{b_m}^{ji}$. Thus, by Proposition IV.2, the times at which $x^{ji}(t^-) = e_m$ and the counter $N_{b_m}^{ji}$ jumps form a renewal sequence with inter-renewal times following the phase-type distribution (ϕ, T) . On the other hand, by Corollary IV.5, these times are exactly the times at which node i attempts to infect node j . Therefore, item c) in Definition III.1 under Assumption III.4 holds true. We can prove the case $t > 0$ in the similar manner and hence omit the proof. $\square$

Before presenting our stability analysis of the GeNeSIS model in the next section, we state another corollary of Lemma IV.4:

Corollary IV.6. *Consider the family of stochastic processes $\xi = \{x^{ji}, y^i\}_{i \in [n], j \in \mathcal{N}_i}$. Then, ξ is a Markov process. Moreover, the states of ξ are all transient except for the absorbing state α at which $x^{ji} = 0$ and $y^i = 0$ for all $i \in [n]$ and $j \in \mathcal{N}_i$. Furthermore, the size of the state space of ξ equals $\prod_{i=1}^n (1 + p^{|\mathcal{N}_i|} q)$, where $|\mathcal{N}_i|$ denotes the size of the set $\mathcal{N}_i$.*

Proof. A careful investigation of the proof of Lemma IV.4 shows the first and the second claims. The third claim is an immediate consequence from the constraint $\mathbb{1}_p^\top x^{ji}(t) = \mathbb{1}_q^\top y^i(t)$. The details are omitted for brevity. $\square$

V. STABILITY ANALYSIS

In this section, we use the results in the previous section to analyze the exponential stability of the GeNeSIS model under Assumption III.4. Specifically, we first present a necessary and sufficient condition for the stability in terms of the eigenvalues of a matrix whose size grows exponentially fast with respect to the relevant parameters. In order to overcome this computational difficulty, we then present an alternative stability condition based on the representation of the GeNeSIS model as the stochastic differential equations (4) and (5).

Throughout this section, we consider the GeNeSIS model with transmission and recovery times following, respectively, phase-type distributions (ϕ, T) and (ψ, R) . The following proposition states a stability condition for the GeNeSIS model, extending the stability analysis on the SIS model in [17]:

Proposition V.1. *Let Π be the infinitesimal generator of the Markov process ξ defined in Corollary IV.6, and let $r < 0$ be the maximum real part of the non-zero eigenvalues of Π . Then, the GeNeSIS model is exponentially mean stable with decay rate λ if and only if $\lambda \leq -r$.*

Proof. Since $\xi(t) \neq \alpha$ if and only if at least one node is infected at time t , we have the inequality $E[z_i(t)] \leq P(\xi(t) \neq \alpha) \leq \sum_{i=1}^n E[z_i(t)]$ for any $i \in [n]$. Therefore, the exponential decay rate of the GeNeSIS model is determined by that of the function $P(\xi(t) \neq \alpha)$ of t . By Corollary IV.6, a basic argument on Markov processes (see [17] for the case of exponential transmission and recover times) shows that *i)* if $\lambda < r$, there exists a constant $C > 0$ such that $P(\xi(t) \neq \alpha) \leq Ce^{-\lambda t}$ for all t and any initial state $\xi(0)$, and *ii)* if $\lambda > r$, there exists an initial state of ξ such that the function $P(\xi(t) \neq \alpha)$ cannot be bounded from above by the exponential function $Ce^{-\lambda t}$ for any value of C . These observations immediately prove the claim of the proposition. $\square$

Although Proposition V.1 gives a necessary and sufficient condition for the exponential mean stability of the GeNeSIS model, the condition is not easily applicable in practice because the size of the matrix Π , i.e., the product $\prod_{i=1}^n (1 + p^{|\mathcal{N}_i|} q)$, grows exponentially fast with respect to the parameters n , p , and q . The following theorem gives a sufficient condition for the exponential stability in terms of the eigenvalues of a matrix whose size grows *linearly* with respect to the parameters in the GeNeSIS model.

Theorem V.2. *Define the $(npq) \times (npq)$ matrix*

$$\mathcal{A} = I_n \otimes (T^\top \oplus R^\top + (\phi b^\top) \otimes I_q) + A \otimes (\phi b^\top) \otimes (\psi \mathbb{1}_q^\top),$$

where A is the adjacency matrix of the graph $\mathcal{G}$ and the vector b is defined in (3). Then, the GeNeSIS model is exponentially stable with decay rate λ if $\lambda \leq -\eta(\mathcal{A})$, where $\eta(\mathcal{A})$ is the spectral abscissa of $\mathcal{A}$.

Proof. Combining equations (4) and (5), we obtain the following $\mathbb{R}^{p+q}$ -valued stochastic differential equation

$$\begin{aligned} d \begin{bmatrix} x^{ji} \\ y^i \end{bmatrix} = & \sum_{m, m'=1}^p \begin{bmatrix} (E_{m'm} - E_{mm}) x^{ji} \\ 0_{nq} \end{bmatrix} dN_{T_{mm'}}^{ji} \\ & + \sum_{m=1}^p \begin{bmatrix} (e_\phi^j e_m^\top - E_{mm}) x^{ji} \\ 0_{nq} \end{bmatrix} dN_{b_m}^{ji} \\ & + \sum_{\ell, \ell'=1}^q \begin{bmatrix} 0_{np} \\ (F_{\ell'\ell} - F_{\ell\ell'}) y^i \end{bmatrix} dN_{R_{\ell\ell'}}^i + \sum_{\ell=1}^q \begin{bmatrix} -x^{ji} y_\ell^i \\ -y^i y_\ell^i \end{bmatrix} dN_{d_\ell}^i \\ & + \sum_{k=1}^n \sum_{m=1}^p a_{ik} \begin{bmatrix} e_\phi^{ji} (1 - \mathbb{1}_q^\top y^i) x_m^{ik} \\ f_\psi^i (1 - \mathbb{1}_q^\top y^i) x_m^{ik} \end{bmatrix} dN_{b_m}^{ik}. \end{aligned}$$

Then, we apply Itô's formula in Lemma II.1 using the function $g(\text{col}(x^{ji}, y^i)) = w^{ji} = x^{ji} \otimes y^i$, to obtain the following

$\mathbb{R}^{pq}$ -valued stochastic differential equation (after tedious, but simple, algebraic manipulations)

$$\begin{aligned}
dw^{ji} &= \left[\sum_{m,m'=1}^p ((E_{m'm} - E_{mm}) \otimes I_q) T_{mm'} \right. \\
&\quad + \sum_{m=1}^p ((E[e_\phi^{ji}] e_m^\top - E_{mm}) \otimes I_q) b_m \\
&\quad + \sum_{\ell,\ell'=1}^q (I_p \otimes (F_{\ell'\ell} - F_{\ell\ell})) R_{\ell\ell'} \Big] \omega_{ji} + \sum_{\ell=1}^q E[-y_\ell^i w^{ji}] d_\ell \\
&\quad + \sum_{k=1}^n \sum_{m=1}^p a_{ik} E[e_\phi^{ji} \otimes f_\psi^i] E[(1 - \mathbb{1}_{pq}^\top w^{ji})(e_m^\top \otimes \mathbb{1}_q^\top) w^{ik}] b_m \\
&= \sum_{m,m'=1}^p ((E_{m'm} - E_{mm}) \otimes I_q) w^{ji} dN_{T_{mm'}}^{ji} \\
&\quad + \sum_{m=1}^p ((e_\phi^{ji} e_m^\top - E_{mm}) \otimes I_q) w^{ji} dN_{b_m}^{ji} \\
&\quad + \sum_{\ell,\ell'=1}^q (I_p \otimes (F_{\ell'\ell} - F_{\ell\ell})) w^{ji} dN_{R_{\ell\ell'}}^i + \sum_{\ell=1}^q (-y_\ell^i w^{ji}) dN_{d_\ell}^i \\
&\quad + \sum_{k=1}^n \sum_{m=1}^p a_{ik} (e_\phi^{ji} \otimes f_\psi^i) (1 - \mathbb{1}_{pq}^\top w^{ji})(e_m^\top \otimes \mathbb{1}_q^\top) w^{ik} dN_{b_m}^{ik}.
\end{aligned} \tag{6}$$

For brevity, we omit the details of this derivation. Define $\omega_{ji}(t) = E[w^{ji}(t)]$ for $t \geq 0$. Then, using Lemma II.2, from (6) we can derive the $\mathbb{R}^{pq}$ -valued differential equation

$$\begin{aligned}
\frac{d\omega_{ji}}{dt} &= \left[(T^\top + B) \otimes I_q + (\phi b^\top - B) \otimes I_q \right. \\
&\quad + \left. (I_p \otimes (R^\top + D)) \right] \omega_{ji} - (I_p \otimes D) \omega_{ji} \\
&\quad + (\phi \otimes \psi)(e_m^\top \otimes \mathbb{1}_q^\top) \sum_{m=1}^p b_m \sum_{k=1}^n (a_{ik} \omega_{ik}) - \varepsilon_{ji}, \tag{7}
\end{aligned}$$

where B and D are the $n \times n$ diagonal matrices having the diagonals $b_1, \dots, b_n$ and $d_1, \dots, d_n$, respectively, and the $\mathbb{R}^{pq}$ -valued function ε_{ji} is defined by $\varepsilon_{ji}(t) = \sum_{k=1}^n \sum_{m=1}^p b_m a_{ik} (\phi \otimes \psi) E[\mathbb{1}_{pq}^\top w^{ji}(t)(e_m^\top \otimes \mathbb{1}_q^\top) w^{ik}(t)]$ for every $t \geq 0$.

Now, for every $i \in [n]$, we take an arbitrary $j_i \in \mathcal{N}_i$. We then define the function ω_i by $\omega_i = \omega_{j_i}$ (i.e., the solution to (7) when $j = j_i$), as well as the $\mathbb{R}^{pq}$ -valued function $\omega = \text{col}(\omega_1, \dots, \omega_n)$. Notice that, from (6), all the stochastic processes in the set $\{w^{ji}\}_{j \in \mathcal{N}_i}$ follow the same stochastic differential equation and, therefore, present the same probability distribution. This implies that $\omega_i = \omega_{j_i}$ for every $j \in \mathcal{N}_i$. Therefore, we can show that $\sum_{k=1}^n a_{ik} \omega_{ik} = \sum_{k=1}^n a_{ik} \omega_k = (A_i \otimes I_{pq}) \omega$, where A_i denotes the i -th row of the adjacency matrix A . Then, from (7), it follows that $d\omega_i/dt = (T^\top \otimes I_q + I_p \otimes R^\top + (\phi b^\top) \otimes I_q) \omega_i + (A_i \otimes (\phi b^\top) \otimes (\psi \mathbb{1}_q^\top)) \omega - \varepsilon_{j_i}$. Defining $\varepsilon = \text{col}(\varepsilon_{j_1}, \dots, \varepsilon_{j_n})$, we obtain the differential equation $d\omega/dt = \mathcal{A}\omega - \varepsilon$, where $\mathcal{A}$ is defined in the theorem.

Since $\mathcal{A}$ is Metzler, we have that $e^{\mathcal{A}t} \geq 0$ for every $t \geq 0$. Also, since both $x^{ji}(t)$ and $y^i(t)$ are nonnegative for all $i \in [n]$, $j \in \mathcal{N}_i$, and $t \geq 0$, we have that $\varepsilon(t) \geq 0$ for every $t \geq 0$. Therefore, it follows that $\omega(t) = e^{\mathcal{A}t} \omega(0) - \int_0^t e^{\mathcal{A}(t-\tau)} \varepsilon(\tau) d\tau \leq e^{\mathcal{A}t} \omega(0)$. This inequality implies that $\omega(t)$ converges to zero as $t \rightarrow \infty$ with a decay rate at least $-\eta(\mathcal{A})$ since $\omega(t) \geq 0$ for all $t \geq 0$. On the other hand, for each $i \in [n]$ and $j \in \mathcal{N}_i$, we can

show that $\mathbb{1}_{pq}^\top w^{ji}(t) = (\mathbb{1}_p^\top x^{ji}(t))(\mathbb{1}_q^\top y^i(t)) = \mathbb{1}_q^\top y^i(t) = z_i(t)$ from Lemma IV.4. Therefore, $E[z_i(t)] = \mathbb{1}_{pq}^\top \omega_{j_i}(t)$, which shows the exponentially fast convergence of $E[z_i(t)]$ towards zero with a decay rate at least $-\eta(\mathcal{A})$. This completes the proof of the theorem. $\square$

Remark V.3. Unlike the necessary and sufficient condition in Proposition V.1, the condition in Theorem V.2 is only sufficient. This conservatism arises from ignoring the higher-order term ε_{ji} in the differential equation (7). Inclusion of these higher-order terms into the analysis would allow us to reduce the conservatism, at the cost of increasing the dimension of the matrix $\mathcal{A}$.

VI. NUMERICAL SIMULATIONS

In this section, we illustrate the effectiveness of our results with numerical simulations in a real social network. We specifically show that modeling non-exponential transmission/recovery times using exponential distributions (i.e., using the standard Markovian model in [17]) can induce significant errors in the computation of the decay rate of the epidemic process. In our simulations, we use the following four distributions to model transmission and recovery times in the GeNeSIS model: (i) the exponential distribution with mean μ (and, hence, variance μ^2); (ii) the log-normal distribution with mean μ and variance μ^2 ; (iii) the log-normal distribution with mean μ and variance $2\mu^2$; and (iv) the log-normal distribution with mean μ and variance $4\mu^2$.

In order to analyze the stability of the GeNeSIS model whose transmission and recovery times follow one of these four distributions, we first approximate the three log-normal distributions in (ii)–(iv) using phase-type distributions (as described in Section III) having $p = 10$ phases. Fig. 1 shows the probability density functions of the (exact) log-normal distributions, as well as the fitted phase-type distributions, when the value of the parameter μ is in the range $[0.5 : 0.1 : 1.5]$.

Using the obtained phase-type distributions, we then apply Theorem V.2 to analyze the stability of the GeNeSIS model when the transmission/recovery times follow one of the four distributions described above. In particular, we compute the exponential decay rate, $-\eta(\mathcal{A})$, for each one of the possible sixteen combinations of transmission/recovery distributions when μ is in the range $[0.5 : 0.05 : 1.5]$. In the subfigures of Fig. 2, we include contour plots of the decay rates for these sixteen cases. We see that, even though the four distributions have the same mean, the resulting GeNeSIS models exhibit different decay rates. In Fig. 2, we observe that for a fixed recovery distribution (i.e., for a fixed column in the table), the colored region (indicated by black solid lines) shrinks as we increase the variance of the log-normal distribution modeling the transmission times (see rows 2 to 4 in Fig. 2). From this observation, we conclude that the ‘heavier’ the tail of the transmission distribution, the more ‘viral’ the spreading process becomes. Similarly, we analyze the impact of the variance of the recovery distribution in the stability of the spread. In Fig. 2, we observe that for a fixed transmission distribution (i.e., for a fixed row in the table), the colored region remains almost unaltered as we increase the variance




Fig. 1. Approximations by phase-type distributions. Solid: Probability density functions of the log-normal distributions. Dashed: Probability density functions of phase-type distributions.




Fig. 2. Lower bound $-\eta(\mathcal{A})$ on the decay rate of generalized SIS model. Horizontal axes: μ . Vertical axes: μ divided by the spectral radius of the graph.

of the distribution modeling the recovery time. However, the decay rate $-\eta(\mathcal{A})$ increases more abruptly inside the blue region as we decrease this variance, as indicated by steeper gradients. In other words, the smaller the variance of the recovery distribution, the faster the spreading process ‘dies out’ in the network.

VII. CONCLUSION

In this paper, we have analyzed the dynamics of a generalized model of spreading over arbitrary networks with phase-type transmission and recovery times. Since phase-type distributions form a dense family in the space of positive-valued distributions, our results allow to theoretically analyze arbitrary transmission and recovery times within an arbitrary

accuracy. In this context, we have derived conditions for a generalized spreading process to converge towards the infection-free equilibrium (i.e., to eradicate the spread) with a given exponential decay rate. In contrast with existing results, we provide an analysis of the *stochastic* spreading dynamics over *arbitrary* networks without relying on mean-field approximations. Our results illustrate that the particular shape of the transmission/recovery distribution heavily influences the decay rate of the convergence towards the infection-free equilibrium. In particular, we observe that the ‘heavier’ the tail of the transmission distribution, the more ‘viral’ the spreading process becomes. Furthermore, the ‘thinner’ the tail of the recovery distribution, the faster the spreading process ‘dies out’ in the network.

REFERENCES

- [1] S. Asmussen, O. Nerman, and M. Olsson, “Fitting phase-type distributions via the EM algorithm,” *Scandinavian Journal of Statistics*, vol. 23, no. 4, pp. 419–441, 1996.
- [2] N. T. J. Bailey, *The Mathematical Theory of Infectious Diseases and its Applications*. Griffin, 1975.
- [3] S. P. Blythe and R. M. Anderson, “Variable infectiousness in HFV transmission models,” *Mathematical Medicine and Biology*, vol. 5, no. 3, pp. 181–200, 1988.
- [4] E. Cator, R. van de Bovenkamp, and P. Van Mieghem, “Susceptible-infected-susceptible epidemics on networks with general infection and cure times,” *Physical Review E*, vol. 87, no. 6, p. 062816, 2013.
- [5] E. Çinlar, *Introduction to Stochastic Processes*. Prentice-Hall, 1975.
- [6] D. R. Cox, “A use of complex probabilities in the theory of stochastic processes,” *Mathematical Proceedings of the Cambridge Philosophical Society*, vol. 51, no. 02, pp. 313–319, 1955.
- [7] A. Ganesh, L. Massoulié, and D. Towsley, “The effect of network topology on the spread of epidemics,” in *24th Annual Joint Conference of the IEEE Computer and Communications Societies*, 2005, pp. 1455–1466.
- [8] R. Horn and C. Johnson, *Matrix Analysis*. Cambridge University Press, 1990.
- [9] H.-H. Jo, J. I. Perotti, K. Kaski, and J. Kertész, “Analytically solvable model of spreading dynamics with non-Poissonian processes,” *Physical Review X*, vol. 4, no. 1, p. 011041, 2014.
- [10] K. Lerman and R. Ghosh, “Information contagion: An empirical study of the spread of news on Digg and Twitter social networks,” in *Fourth International AAAI Conference on Weblogs and Social Media*, 2010, pp. 90–97.
- [11] C. Nowzari, V. M. Preciado, and G. J. Pappas, “Analysis and control of epidemics: A survey of spreading processes on complex networks,” *IEEE Control Systems*, vol. 36, no. 1, pp. 26–46, 2016.
- [12] M. Ogura and V. M. Preciado, “Stability of spreading processes over time-varying large-scale networks,” *IEEE Transactions on Network Science and Engineering*, vol. 3, no. 1, pp. 44–57, 2016.
- [13] —, “Epidemic processes over adaptive state-dependent networks,” *Physical Review E*, vol. 93, p. 062316, 2016.
- [14] B. Øksendal, *Stochastic Differential Equations*. Springer, 2003.
- [15] L. Pellis, T. House, and M. J. Keeling, “Exact and approximate moment closures for non-Markovian network epidemics,” *Journal of Theoretical Biology*, vol. 382, pp. 160–177, 2015.
- [16] T. Suess, U. Buchholz, S. Dupke, R. Grunow, M. An der Heiden, A. Heider, B. Biere, B. Schweiger, W. Haas, and G. Krause, “Shedding and transmission of novel influenza virus A/H1N1 infection in households-Germany, 2009,” *American Journal of Epidemiology*, vol. 171, no. 11, pp. 1157–1164, 2010.
- [17] P. Van Mieghem, J. Omic, and R. Kooij, “Virus spread in networks,” *IEEE/ACM Transactions on Networking*, vol. 17, no. 1, pp. 1–14, 2009.
- [18] P. Van Mieghem and R. van de Bovenkamp, “Non-Markovian infection spread dramatically alters the susceptible-infected-susceptible epidemic threshold in networks,” *Physical Review Letters*, vol. 110, no. 10, p. 108701, 2013.